

	<p>Council</p> <p><i>Genetic and Other Innovative Strategies to Reduce Sea Lice</i></p>	<p>CNL(21)47</p> <p>Agenda Item: 5(a)</p>
---	--	---

Genetic and Other Innovative Strategies to Reduce Sea Lice

Michael Pietrak and Brian Peterson

U.S. Department of Agriculture, Agricultural Research Service, National Cold Water Marine Aquaculture Research Center, 25 Salmon Farm Rd, Franklin, ME 04634 USA

The United States Department of Agriculture-Agricultural Research Service (USDA-ARS) National Cold Water Marine Aquaculture Center (NCWMAC) in Franklin, ME has been supporting the U.S. coldwater marine aquaculture industry since 2003 by developing genetically improved North American (NA) Atlantic salmon and conducting research to address industry issues. In the past six years the NCWMAC has begun two programs to help manage sea lice on marine net pen farms. The first program incorporates sea lice resistance into the existing selective breeding program and the second is the development of lumpfish culture techniques and broodstock for use as cleaner fish in marine net pens. Both selective breeding and the use of cleaner fish are management options that can be used by the domestic industry as part of their existing integrated pest management programs (IPMP).

Breeding Program

The NCWMAC utilizes the St. John River (SJR) strain because of fast growth, certification of NA origin, and widespread utilization by industry. Since 2015, the primary objectives of the salmon selection program have been to improve commercially important traits including: carcass weight, fillet color, fat content, and sea lice resistance. Recently, the program started to evaluate and validate the usefulness of incorporating genomic based selection techniques into the salmon breeding program. Culture of Atlantic salmon in the breeding program and research facility is based on life stage and separation of year class (Figure 1). An incubation system is used for eggs and newly hatched fry before first feeding, parr tanks are used for first feeding fry to 30-40 g salmon, the smolt tanks are for 30-100 g pit tagged salmon, on-grow tanks are for 100 g to 1.0 kg salmon in their second year, 3-yr old broodstock tanks are for 1.0 kg to 3.0 kg salmon, and four 4-yr old broodstock tanks are for growing salmon 3.0 to 8.0 kg when they will be spawned. Up to 234 families of Atlantic salmon can be cultured in the breeding nucleus and approximately 250 eggs are saved from each family mating and reared. The remaining eggs are distributed to industry and used as their next generation of brood to provide eggs for commercial production. Of the fish from the 250 eggs that remain in the program, typically, 30-40 smolts per family are saved as the nucleus of fish (broodstock) for the breeding program and cultured in biosecure tanks at the NCWMAC. An additional 30-40 smolts per family are cultured and transferred to an industry collaborator for stocking into net pens for growth performance evaluations. Additional smolts are also kept for various studies involving measurement of traits such as sea lice resistance. Selection of 4-year old fish for spawning occurs when fish are moved from 3-yr old broodstock tanks into the 4-yr broodstock system prior to the spawning season.

Sea Lice Selection

Since 2015, the NCWMAC has evaluated three of the four year classes per generation and continues to evaluate and select the offspring of those year classes. The program utilizes a challenge model based on multiple small challenges to evaluate 70-120 families for lice resistance. Approximately 16 pit tagged fish from each of these families are set aside for

challenge trials. One fish from each family is stocked into a 1 m³ tank, setting up a series of 16 replicate tanks.

Sea lice egg strings are collected from commercial farms and then brought back to be cultured in the laboratory to the infective copepodid stage. Challenge fish are infested via a 4-hour bath challenge as 2-4 day old copepodids are available. Lice are allowed to develop to the 2nd chalimus stage, approximately 10-14 days post infection (dpi). Fish are euthanized by an overdose of MS-222 in saltwater and then transferred to a freshwater staining bath of neutral red for 20 min before being individually bagged. Fish are kept on ice until the lice are counted, fish weighed, fork length measured, and fin clips are preserved in 95% Ethanol. Lice counts are standardized by lice density (LD; Gjerde *et al.* 2011) and lice per cm² of surface area. Surface area is calculated from the model in Frederick *et al.* (2017). Estimated breeding values for each family are generated using the program MTDREL.

The evaluations to date have revealed heritability values within our population of between 0.20 and 0.31 that are consistent with those published in the literature. Looking at the average lice densities for each family in each year reveals a similar pattern with a few families showing resistance and a gradual increase towards highly susceptible families (Figure 2). Efforts to date have focused on utilizing the most resistant families to select a few highly resistant families, while simultaneously, ensuring that most families which are selected based on growth are families not within the most susceptible 30% of families.

Genomic Based Selection

Genomic based selection has made significant advances over the past 20 years since the development of the use of genome-wide dense marker maps (Meuwissen *et al.* 2001). In combination with the reduction in genotyping costs this new technique has allowed for improved selection accuracy. This is accomplished by using a combination of traditional phenotypic based estimated breeding values (EBV) and genomic data to generate genomic estimated breeding values (gEBV) which account for within family variation in addition to between family variation. The techniques were pioneered in dairy and beef cattle and other major terrestrial livestock, but recently are more widely being used in fish for multi-locus traits such as growth and disease resistance to some pathogens (Vallejo *et al.* 2017; Garcia *et al.* 2018).

Over the past several years the program has been collaborating with USDA-ARS scientists at the National Center for Cool and Cold Water Aquaculture (NCCCWA) in Leetown, WV. These efforts have included constructing a reference genome for North American Atlantic salmon of the SJR strain and developing a publicly available 50K single nucleotide polymorphism (SNP) chip for genotyping salmon of NA origin (Gao *et al.* 2020). The goal is to implement a genomic selection program for sea lice in November of 2021. This represents a significant step forward in selective breeding for sea lice resistance.

Lumpfish

Cleaner fish have been used in European salmon farms for over 30 years (Pike 1989). However, the wrasse species that have been primarily used are not native to the western North Atlantic Ocean. More recently the lumpfish (*Cyclopterus lumpus* L.) has become a popular option for use as a cleaner fish. Lumpfish are native to the eastern coast of North America (Cox and Anderson 1918). As a result, the use of cleaner fish has only been adopted within the past 5 years or so with the commercial industry on the North American side of the Atlantic Ocean. Despite, the disparate adoptions in the use of cleaner fish between the two sides of the Atlantic, the recent surge in use of lumpfish is common to both and the development of a lumpfish farming industry is rapidly developing in both locations. This rapid development presents a

range of research potential and knowledge gaps for the species (Brooker *et al.* 2018; Powell *et al.* 2018).

The wide range of research coming out on lumpfish, can be grouped into four broad areas. The first is the development of captive reared broodstock populations. The reliance on wild caught animals to provide eggs has been identified in several key reviews as a critical knowledge gap (Brooker *et al.* 2018; Powell *et al.* 2018). However, this situation is rapidly evolving as development of captive brood is constantly occurring in major lumpfish rearing countries. This research area is evolving to incorporate potential selective breeding, manipulation of spawning timing and improving reproduction (Imsland *et al.* 2016a; Imsland *et al.* 2019). This is one of the primary areas of focus for our lumpfish program. In collaboration with the University of Maine, current efforts are focused on development of captive brood populations derived from local populations. Likewise, the U.S. Lumpfish Consortium spearheaded by NCWMAC, University of New Hampshire, the University of Maine and industry partners are also investigating the manipulation of spawning and interested in the potential for selective breeding.

The second, broad area of research involves optimizing the effective husbandry of all stages of lumpfish care both in hatcheries and in sea cages. There is a wide range of issues and topics being explored under this area. Work in this area is coming out in both peer reviewed and trade literature but is also being conducted within facilities and companies on a private basis. On the hatchery side, research into optimum water quality, temperature, nutrition and other husbandry issues are being conducted (D'Arcy *et al.* 2020; Dahle *et al.* 2020; Mortensen *et al.* 2020; Pountney *et al.* 2020). Optimizing hatchery husbandry including the maintenance and nutrition of captive broodstock populations is the other primary focus of the NCWMAC in collaboration with the University of New Hampshire. On the net pen side nutrition, lumpfish recapture and housing are areas of interest (Imsland *et al.* 2015a; Imsland *et al.* 2018a; Imsland *et al.* 2018b; Imsland *et al.* 2020) Feeding techniques, such as, use of gel blocks or submersible feeders to keep lumpfish from eating salmon feed are also being examined. One of the more interesting trends that has occurred with the development of lumpfish culture is not only is there interest in optimized husbandry as demonstrated above, but there is interest in animal welfare research and how to improve this research simultaneously (Gutierrez Rabadan *et al.* 2020).

The third major area of research revolves around disease. The development of any new aquaculture species is accompanied by many new areas for disease research as it is difficult to study disease in wild populations. As might be expected, new occurrences of bacterial, viral, parasitic and mycotic pathogens are being observed (Einarsdottir *et al.* 2018; Ellul *et al.* 2018; Rouleau *et al.* 2018; Scholz *et al.* 2018; Pietrak and Rosser 2019; Stagg *et al.* 2020). Research into lumpfish immunity, pathogen treatment and other areas to improve overall health is also rapidly developing (Erkinharju *et al.* 2018; Pietrak and Backman 2018; Haugland *et al.* 2019; Jacobsen *et al.* 2019; Eggestøl *et al.* 2020; Klakegg *et al.* 2020). One new area of health research that is and will become more prominent is the ecology of pathogens in these polyculture systems (Murray 2017). Large scale western aquaculture has traditionally been a monospecies approach and the benefits of polyculture systems has focused on multi-trophic systems rather than multiple fish species. The increased co-culture of cleaner fish can complicate disease management on farms given the ability of some pathogens or strains to remain asymptomatic in one species cultured, but potentially cause outbreaks the other.

Finally, the last major area of research is arguably the most important and that is documenting and optimizing the efficacy of using lumpfish as cleaner fish. Early efforts focused on demonstrating the effectiveness of lumpfish by comparing sea lice levels in cages with and without lumpfish, and with varying densities of lumpfish, and other factors (Imsland *et al.*

2014a; Imsland *et al.* 2014b; Imsland *et al.* 2015; Imsland *et al.* 2016b; Imsland *et al.* 2018c). However, while there is a consensus that lumpfish can help manage sea lice populations, it has been observed that not all lumpfish exhibit cleaning behavior. A pressing topic is understanding what genetic, environmental, seasonal, or life history factors are influencing cleaning behavior (Imsland *et al.* 2016a; Imsland *et al.* 2016b; Eliassen *et al.* 2018; Leclercq *et al.* 2018). By starting to understand the various factors influencing cleaning behavior, then studies can start to help refine and optimize the most appropriate situations for utilizing lumpfish (McEwan *et al.* 2019).

Summary

Significant advances are being made in the development of selective breeding for sea lice resistance and in the use of cleaner fish, especially lumpfish. However, neither of these technologies likely represents a silver bullet to managing sea lice populations on either a local or global scale. Both technologies can be important tools and components in effective IPMP as salmon farmers move from IPMP that primarily use drug-based management strategies to IPMP that use non-drug based strategies. It is important to remember that no specific system of control strategies will fit all regions, water bodies or even companies. Rather systems need to be individually tailored to each farm. With the proliferation of non-drug management tools in IPMP, it is critical that more research be focused on rigorously determining what strategies and technologies are effective at the local level and defining the economic costs of each strategy.




Figure 1. Overview of the 4-year cycle for the breeding program at the U.S. Department of Agriculture’s National Cold Water Marine Aquaculture Center.




Figure 2. Typical graph illustrating average lice density (LDTotal) per family with 95% confidence intervals. Families exhibiting lower lice density are more resistant.

References

- Brooker, A.J., Papadopoulou, A., Gutierrez, C., Rey, S., Davie, A. and Migaud, H. 2018. Sustainable production and use of cleaner fish for the biological control of sea lice: recent advances and current challenges. *Veterinary Record*, 183, 383.
- D'Arcy, J., Kelly, S., McDermott, T., Hyland, J., Jackson, D. and Bolton-Warberg, M. 2020. Assessment of PIT tag retention, growth and post-tagging survival in juvenile lumpfish, *Cyclopterus lumpus*. *Animal Biotelemetry*, 8, 1.
- Dahle, S.W., Bakke, I., Birkeland, M., Nordøy, K., Dalum, A.S. and Attramadal, K.J.K. 2020. Production of lumpfish (*Cyclopterus lumpus* L.) in RAS with distinct water treatments: Effects on fish survival, growth, gill health and microbial communities in rearing water and biofilm. *Aquaculture*, 522, 735097.
- Eggestøl, H.Ø., Lunde, H.S. and Haugland, G.T. 2020. The proinflammatory cytokines TNF- α and IL-6 in lumpfish (*Cyclopterus lumpus* L.) -identification, molecular characterization, phylogeny and gene expression analyses. *Developmental & Comparative Immunology*, 105, 103608.
- Einarsdottir, T., Sigurdardottir, H., Bjornsdottir, T.S. and Einarsdottir, E. 2018. *Moritella viscosa* in lumpfish (*Cyclopterus lumpus*) and Atlantic salmon (*Salmo salar*). *Journal of Fish Diseases*.
- Eliassen, K., Danielsen, E., Johannesen, Á., Joensen, L.L. and Patursson, E.J. 2018. The cleaning efficacy of lumpfish (*Cyclopterus lumpus* L.) in Faroese salmon (*Salmo salar* L.) farming pens in relation to lumpfish size and seasonality. *Aquaculture*, 488, 61-65.
- Ellul, R.M., Walde, C., Haugland, G.T., Wergeland, H. and Ronneseth, A. 2018. Pathogenicity of *Pasteurella* sp. in lumpsuckers (*Cyclopterus lumpus* L.). *Journal of Fish Diseases*.

- Erkinharju, T., Dalmo, R.A., Vågsnes, Ø., Hordvik, I. and Seternes, T. 2018. Vaccination of Atlantic lumpfish (*Cyclopterus lumpus* L.) at a low temperature leads to a low antibody response against *Aeromonas salmonicida*. *Journal of Fish Diseases*, 41.
- Frederick, C., Brady, D.C. and Bricknell, I. 2017. Landing strips: Model development for estimating body surface area of farmed Atlantic salmon (*Salmo salar*). *Aquaculture*, 473, 299-302.
- Gao, G., Pietrak, M.R., Burr, G.S., Rexroad, C.E., Peterson, B.C. and Palti, Y. 2020. A New Single Nucleotide Polymorphism Database for North American Atlantic Salmon Generated Through Whole Genome Resequencing. *Frontiers in Genetics*, 11,
- Garcia, A.L.S., Bosworth, B., Waldbieser, G., Misztal, I., Tsuruta, S. and Lourenco, D.A.L. 2018. Development of genomic predictions for harvest and carcass weight in channel catfish. *Genetics Selection Evolution*, 50, 66.
- Gjerde, B., Odegard, J. and Thorland, I. 2011. Estimates of genetic variation in the susceptibility of Atlantic salmon (*Salmo salar*) to the salmon louse *Lepeophtheirus salmonis*. *Aquaculture*, 314, 66-72.
- Gutierrez Rabadan, C., Spreadbury, C., Consuegra, S. and Garcia de Leaniz, C. 2020. Development, validation and testing of an Operational Welfare Score Index for farmed lumpfish *Cyclopterus lumpus* L. *Aquaculture*.
- Haugland, G.T., Kverme, K.O., Hannisdal, R., Kallekleiv, M., Colquhoun, D.J., Lunestad, B.T., Wergeland, H.I., *et al.* 2019. Pharmacokinetic Data Show That Oxolinic Acid and Flumequine Are Absorbed and Excreted Rapidly From Plasma and Tissues of Lumpfish. *Frontiers in Veterinary Science*, 6, 394.
- Imslund, A., Reynolds, P., Eliassen, G., Hangstad, T., Nytrø, A., Foss, A., Vikingstad, E., *et al.* 2015a. Assessment of suitable substrates for lumpfish in sea pens. *Aquaculture International*, 23, 639-645.
- Imslund, A.K., Hangstad, T.A., Jonassen, T.M., Stefansson, S.O., Nilsen, T.O., Hovgaard, P., Elvegård, T.A., *et al.* 2019. The use of photoperiods to provide year round spawning in lumpfish *Cyclopterus lumpus*. *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology*, 228, 62-70.
- Imslund, A.K., Reynolds, P., Eliassen, G., Hangstad, T.A., Foss, A., Vikingstad, E. and Elvegård, T.A. 2014a. The use of lumpfish (*Cyclopterus lumpus* L.) to control sea lice (*Lepeophtheirus salmonis* Krøyer) infestations in intensively farmed Atlantic salmon (*Salmo salar* L.). *Aquaculture*, 424-425, 18-23.
- Imslund, A.K., Reynolds, P., Eliassen, G., Hangstad, T.A., Nytrø, A.V., Foss, A., Vikingstad, E., *et al.* 2015b. Feeding preferences of lumpfish (*Cyclopterus lumpus* L.) maintained in open net-pens with Atlantic salmon (*Salmo salar* L.). *Aquaculture*, 436, 47-51.
- Imslund, A.K., Reynolds, P., Eliassen, G., Hangstad, T.A., Nytrø, A.V., Foss, A., Vikingstad, E., *et al.* 2014b. Assessment of growth and sea lice infection levels in Atlantic salmon stocked in small-scale cages with lumpfish. *Aquaculture*, 433, 137-142.
- Imslund, A.K., Reynolds, P., Eliassen, G., Mortensen, A., Hansen, Ø.J., Puvanendran, V., Hangstad, T.A., *et al.* 2016a. Is cleaning behaviour in lumpfish (*Cyclopterus lumpus*) parentally controlled? *Aquaculture*, 459, 156-165.

- Imslund, A.K., Reynolds, P., Hangstad, T.A., Jónsdóttir, Ó.D.B., Noble, T., Wilson, M., Mackie, J.A., *et al.* 2018a. Feeding behaviour and growth of lumpfish (*Cyclopterus lumpus* L.) fed with feed blocks. *Aquaculture Research*.
- Imslund, A.K., Reynolds, P., Nytrø, A.V., Eliassen, G., Hangstad, T.A., Jónsdóttir, Ó.D.B., Emaus, P.-A., *et al.* 2016b. Effects of lumpfish size on foraging behaviour and co-existence with sea lice infected Atlantic salmon in sea cages. *Aquaculture*, 465, 19-27.
- Imslund, A.K.D., Reynolds, P., Jonassen, T.M., Hangstad, T.A., Elvegård, T.A., Urskog, T.C. and Mikalsen, B. 2018b. Effects of three commercial diets on growth, cataract development and histopathology of lumpfish (*Cyclopterus lumpus* L.). *Aquaculture Research*, 49, 3131-3141.
- Imslund, A.K.D., Hanssen, A., Nytro, A.V., Reynolds, P., Jonassen, T.M., Hangstad, T.A., Elvegard, T.A., *et al.* 2018c. It works! Lumpfish can significantly lower sea lice infestation in large-scale salmon farming. *Biology Open*, 7.
- Imslund, A.K.D., Reynolds, P., Lorentzen, M., Eilertsen, R.A., Micallef, G. and Tvenning, R. 2020. Improving survival and health of lumpfish (*Cyclopterus lumpus* L.) by the use of feed blocks and operational welfare indicators (OWIs) in commercial Atlantic salmon cages. *Aquaculture*, 527, 735476.
- Jacobsen, J.V., Steen, K. and Nilssen, K.J. 2019. Anaesthetic efficacy of Aqui-S, Benzoak, and MS-222 on lumpfish (*Cyclopterus lumpus*) fries. Impact from temperature, salinity, and fasting. *PLOS One*, 14, e0211080.
- Klakegg, Ø., Myhren, S., Juell, R.A., Aase, M., Saloniis, K. and Sørum, H. 2020. Improved health and better survival of farmed lumpfish (*Cyclopterus lumpus*) after a probiotic bath with two probiotic strains of *Aliivibrio*. *Aquaculture*, 518, 734810.
- Leclercq, E., Zerafa, B., Brooker, A.J., Davie, A. and Migaud, H. 2018. Application of passive-acoustic telemetry to explore the behaviour of ballan wrasse (*Labrus bergylta*) and lumpfish (*Cyclopterus lumpus*) in commercial Scottish salmon sea-pens. *Aquaculture*, 495, 1-12.
- McEwan, G.F., Groner, M.L., Cohen, A.A.B., Imslund, A.K.D. and Revie, C.W. 2019. Modelling sea lice control by lumpfish on Atlantic salmon farms: interactions with mate limitation, temperature and treatment rules. *Diseases of Aquatic Organisms*, 133, 69-82.
- Meuwissen, T.H.E., Hayes, B.J. and Goddard, M.E. 2001. Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps. *Genetics*, 157, 1819-1829.
- Mortensen, A., Johansen, R.B., Hansen, Ø.J. and Puvanendran, V. 2020. Temperature preference of juvenile lumpfish (*Cyclopterus lumpus*) originating from the southern and northern parts of Norway. *Journal of Thermal Biology*, 89, 102562.
- Murray, A.G. 2017. A model of the process of spillover and adaption leading to potential emergence of disease in salmon held with cleaner fish used to control lice. *Aquaculture*, 473, 283-290.
- Pietrak, M. and Backman, S. 2018. Treatment of lumpfish (*Cyclopterus lumpus* L.) infected with *Gyrodactylus cyclopteri* (Scyborskaya 1948). *Journal of Fish Diseases*, 41, 721-723.
- Pietrak, M. and Rosser, T.G. 2019. Morphologic and molecular characterization of *Gyrodactylus cyclopteri* Scyborskaja, 1948, from *Cyclopterus lumpus* L., 1758. *Parasitology Research*,
- Pike, A.W. 1989. Sea lice — Major pathogens of farmed atlantic salmon. *Parasitology Today*, 5, 291-297.

- Pountney, S.M., Lein, I., Migaud, H. and Davie, A. 2020. High temperature is detrimental to captive lumpfish (*Cyclopterus lumpus* L) reproductive performance. *Aquaculture*, 522, 735121.
- Powell, A., Treasurer, J.W., Pooley, C.L., Keay, A.J., Lloyd, R., Imsland, A.K. and Garcia de Leaniz, C. 2018. Use of lumpfish for sea-lice control in salmon farming: challenges and opportunities. *Reviews in Aquaculture*, 10, 683-702.
- Rouleau, F.D., Vincent, A.T. and Charette, S.J. 2018. Genomic and phenotypic characterization of an atypical *Aeromonas salmonicida* strain isolated from a lumpfish and producing unusual granular structures. *Journal of Fish Diseases*, 41, 673-681.
- Scholz, F., Ruane, N., Marcos-López, M., Mitchell, S., Bolton-Warberg, M., O'Connor, I., Mirimin, L., *et al.* 2018. Systemic mycoses in lumpfish (*Cyclopterus lumpus* L.) in Ireland: aetiology and clinical presentation. *Bulletin of the European Association of Fish Pathologists*, 35, 202-212.
- Stagg, H.E.B., Guðmundsdóttir, S., Vendramin, N., Ruane, N.M., Sigurðardóttir, H., Christiansen, D.H., Cuenca, A., *et al.* 2020. Characterization of ranaviruses isolated from lumpfish *Cyclopterus lumpus* L. in the North Atlantic area: proposal for a new ranavirus species (European North Atlantic Ranavirus). *Journal of General Virology*.
- Vallejo, R.L., Leeds, T.D., Gao, G., Parsons, J.E., Martin, K.E., Evenhuis, J.P., Fragomeni, B.O., *et al.* 2017. Genomic selection models double the accuracy of predicted breeding values for bacterial cold water disease resistance compared to a traditional pedigree-based model in rainbow trout aquaculture. *Genetics Selection Evolution*, 49, 17.