

**Council**

**CNL(04)31**

*Predator-related Mortality  
(tabled by the United States)*

***Predator-related Mortality***  
***(tabled by the United States)***

**Introduction**

Predation has long been recognized as a potential factor that limits the abundance of Atlantic salmon (*Salmo salar*) populations throughout their range. In freshwater environments, predation has been relatively well studied and in some cases these studies have led to a more quantitative understanding of the salmon's role in aquatic food webs (e.g., Larsson 1985, van den Ende 1993). However in marine environments, this level of understanding has not been achieved. This understanding is lacking because of the varying spatial scales at which populations can be studied in the open ocean, the myriad of potential predators that salmon encounter, and the relatively small role that salmon play in marine food webs (Cairns 2001a). A recent review of the potential causes for declines in salmon abundance (Cairns 2001b) underscores the importance of considering marine predation with five of the leading 12 hypotheses being related to predation. The problems of understanding the role of predation in structuring US salmon stocks are exacerbated by critically low abundance levels. Abundance of remnant US stocks are presently so low that the National Marine Fisheries Service and the US Fish and Wildlife Service recently determined that the Gulf of Maine Distinct Population Segment (DPS) is in danger of extinction throughout its range (65 Fed Reg 69459). The purposes of this paper are to (1) summarize recent information regarding the impact of marine predators on US Atlantic salmon stocks and (2) to describe management measures and on-going research in relation to predator-related mortality.

**The Impact of Marine Predators on US Stocks of Atlantic Salmon**

Recent research has advanced our understanding of the impacts of marine predation on US stocks of Atlantic salmon. These advances include both species-specific investigations as well as investigations that clarify the mechanisms for particularly high predation at various life stages, especially as smolts and post-smolts transition to life at sea. We will first consider the recent literature pertaining to species-specific interactions.

*Seabirds*

Northern gannets (*Morus bassanus*) are the largest seabirds in the north Atlantic that can prey on post-smolts throughout their first year at sea (Cairns 2001a). Montevecchi et al. (2002) examined the diets of northern gannets on Funk Island, Newfoundland for 24 years. Atlantic salmon post-smolts comprised between 0 and 6.37% of the northern gannets' diet. Atlantic salmon consumption by northern gannets varied greatly across years but consumption was highest during the 1990s. Overall, northern gannets consumed an estimated 0.22% of North American post-smolt biomass in August from 1977 to 1989 and 2.70% of North American post-smolt biomass in August from 1990 to 2000. The highest consumption of Atlantic salmon occurred in 1993 when northern gannets consumed an estimated 43.4 tons of salmon in August alone. The higher post-smolt predation rates in the 1990s appear to be associated with colder surface water, though many other factors may have affected this shift (e.g., the collapse of ground fish stocks). Tags of salmon from US stocks (Penobscot River, Maine) were found in this colony. The extent of predation by northern gannets specific to US stocks

remains unclear though it is likely to be roughly equal to the overall proportion of the US stock in the Northwest Atlantic mixed stock complex.

Unlike northern gannets, double-crested cormorants (*Phalacrocorax auritus*) have long been recognized as important predators of smolts and post-smolts (Meister and Gramlich 1967). Breeding pairs of cormorants in Maine have increased by 87% between 1977 and 1985 to around 29,000 pairs and this population been stable since the early 1990's (Baum 1997, Anthony 1994). Blackwell et al. (1997) demonstrated that smolts were among the most frequent prey items of double-crested cormorants in the Penobscot River, Maine during the smolt emigration period. Observations by Blackwell et al. (1997) and Blackwell and Krohn (1997) suggest that cormorants prefer to forage near dams where smolts are delayed above the dam and smolts that have gone through the dam are disoriented. These mechanisms may lead to the high levels of predation observed by Moring et al. (1999). Recent studies in eastern Canada also underscore the importance of predation by cormorants on emigrating smolts. Smolts comprised 3.3% of cormorants' diet in the Maritime Provinces during the smolt run (Cairns 1998). This study was not targeted at the major salmon rivers and instead represents an overall assessment of cormorant diet throughout the study area. Thus, overall predation intensity by cormorants is still unknown (Cairns 2001a).

Vulnerability windows have recently been established for most of the major seabird predators of salmon (Cairns and Reddin 2000). Smolts and post-smolts are vulnerable to seabird predation as soon as they leave the river until they attain a critical size at which they are no longer vulnerable to a specific predator. Post-smolts out-grow northern fulmars (*Fulmarus glacialis*), black-legged kittiwakes (*Rissa tridactyla*), and common murrelets (*Uria aalge*) by early July; shearwaters (*Puffinus* spp.) by late July; and gulls (*Larus* spp.) by early September (Cairns and Reddin 2000). This review did not include great cormorants (*Phalacrocorax carbo*) or double-crested cormorants because they are inshore predators and salmon are only available as prey as they migrate through estuaries and near-shore areas. Northern gannets were also not included in this review. However, the vulnerability window to northern gannet predation is known to be considerably wider than other seabirds because of specialized feeding adaptations that allow them to feed on medium-size fish (Cairns and Reddin 2000).

### *Marine Mammals*

Direct evidence of salmon predation by seals in the northwest Atlantic remains sparse. One harbor seal (*Phoca vitulina*), two grey seals (*Halichoerus grypus*), and two harp seals (*Phoca groenlandica*) have been documented with wild salmon remains or tags in their stomachs (Cairns 2001a). Seal predation is better known in the northeastern Atlantic, especially the UK, where seals affect commercial fisheries (Harwood 1984, Gulland 1987). Predation on free-swimming salmon is not as well documented. Most records of seal predation occur near salmon nets (Pierce et al. 1991).

Harbor seals are of particular concern in Maine, USA where seal bites on returning wild salmon have increased during the last decade (Baum 1997). Harbor seal populations in Maine have increased sharply over the last two decades (Gilbert 2003, pers. comm.). This increase clearly coincides with more bite marks on salmon, however actual consumption of wild Maine salmon by harbor seals has never been recorded.

Harbor seals also pose an indirect threat to endangered salmon by damaging aquaculture net pens. Not only are seal attacks on the pens a significant cause of predation, they also damage

nets. These attacks can lead to large escapes of domesticated salmon which may lead to disruption of wild salmon redds, competition for food and habitat, interbreeding with wild salmon, disease transmission, and benthic habitat degradation (Windsor and Hutchinson 1990, Saunders 1991, Webb et al. 1993, Youngson et al. 1993, Clifford et al. 1998).

Another potential salmon predator is the grey seal which ranges across the north Atlantic to the southern Gulf of Maine, inhabiting both coastal and offshore waters. Predation on salmon in the Gulf of Maine by grey seals has not been documented. However, predation by grey seals has been documented in the northwest Atlantic on two occasions (Cairns and Reddin 2000). Salmon remains have been found in the stomachs of grey seals from the Gulf of St. Lawrence, although no salmon remains were found in the stomachs of grey seals from the Scotian Shelf (Anthony 1994). All other accounts of grey seals feeding on salmon are from the northeastern Atlantic. Salmon predation by seals has been documented in many studies conducted in the UK though many incidences of predation are likely net-caught fish (Boyle et al. 1990).

Harp seals are abundant in the north Atlantic and usually inhabit pack ice (Lawson et al. 1998). Harp seal populations in the north Atlantic have increased during the last decade, with breeding populations in Newfoundland, the Gulf of St. Lawrence, and Greenland (Hammill and Stenson 2000). Harp seals feed selectively on capelin (*Mallotus villosus*) and arctic cod (*Boreogadus saida*) but will prey on other available fishes (Lawson et al. 1998). Hammill and Stenson (2000) estimated that harp seals consumed 82% (3.3 million t) of all food eaten by seals in eastern Canadian waters in 1996. Thus if salmon predation is proportional to overall consumption, harp seals may be important salmon predators even though predation on wild salmon in the northwest Atlantic has only been recorded twice (Cairns and Reddin 2000). Further, the abundance of harp seals appears to be inversely correlated with salmon recruitment in the north Atlantic (Amiro 1998). However, harp seals may simply be a surrogate for some other predator or environmental condition that reduces salmon recruitment.

Overall, seal predation is likely to have risen substantially over the last decade concurrent with increases in overall seal abundance in the northwest Atlantic. Hammill and Stenson (2000) estimated that overall consumption of salmon by harp, hooded (*Cystophora cristata*), grey, and harbor seals steadily increased from 2,467 t in 1990 to 3,229 t in 1996. However, the vast majority (93% in 1996) of all salmon predation by seals is thought to occur in NAFO areas 4R and 4S (i.e., the western side of Newfoundland). US salmon stocks are thought to migrate through and over-winter on the eastern side of Newfoundland (Meister 1984, Baum 1997). Though this migration pattern is not known with certainty, Hammill and Stenson's (2000) results suggest that post-smolts migrating through and over-wintering on the eastern side of Newfoundland may have considerably lower predation risks than those on the western side of Newfoundland. Thus, US salmon stocks may be subject to substantially lower seal predation risks than many Canadian stocks. However, these consumption estimates, and subsequent predation rates, are based on the few records of salmon predation available.

As with seabirds, vulnerability windows were recently established for the major seal predators of salmon (Cairns and Reddin 2000). Vulnerability windows for hooded, harbor, and grey seals are quite variable and depend largely on spatial co-occurrence. Vulnerability for harp seals is more straight-forward with both post-smolts and 1SW salmon being vulnerable throughout the year.

The extent of predation by other marine mammals is still poorly understood. The only indication that cetaceans may prey on salmon is from a single salmonid otolith found in a harbor porpoise (*Phocoena phocoena*) stomach (Cairns 2001b), though Thompson and Mackay (1999) suggest that many of the marks attributed to seals may have been inflicted by odontocete cetaceans instead. Without meaningful consumption estimates, it is currently impossible to assess the impacts of cetacean predation. However, the distribution of several odontocete cetaceans clearly overlaps the distribution of Atlantic salmon. Furthermore, Pacific salmon (*Oncorhynchus* spp.) have been reported from the stomachs of nine species of cetaceans (Fiscus 1980). Thus, cetaceans may harvest a substantial portion of Atlantic salmon biomass even if salmon are a small portion of their diet (Cairns 2001b).

### *Fish*

Striped bass (*Morone saxatilis*) are presumed to be important predators of smolts and post-smolts as they migrate through nearshore waters. This presumption has recently gained credibility by two studies that documented the intensity (Blackwell and Juanes 1998) and the spatial variability (Beland et al. 2001) of striped bass predation on Atlantic salmon. Blackwell and Juanes (1998) documented striped bass preying on salmon smolts in the tailrace of the Essex Dam on the Merrimack River, Massachusetts, USA. Smolts represented approximately 80% of prey recovered. This was during the peak smolts emigration period (May 6 to May 28) and most smolts were hatchery-origin fish either stocked as fry or smolts. In addition, Schulze (1996) demonstrated that the spatial and temporal distributions of striped bass and emigrating smolts in the Connecticut River estuary overlap substantially and further concluded that only early-migrating smolts would not be subject to striped bass predation. Striped bass predation has also been documented in the Narraguagas River (Beland et al. 2001) which is one of the rivers containing an endangered population of Atlantic salmon in eastern Maine, USA (65 Fed Reg 69459). Though the extent of striped bass predation on endangered salmon is still unclear, the risk of population effects upon salmon is presently high given low smolt production (Beland et al. 2001) and high striped bass abundance (Field 1997).

### *Underlying Mechanisms*

Several of the underlying mechanisms for the apparently high marine predation observed by many researchers are becoming clear. Some of these mechanisms include water quality perturbations and decreased abundance of other anadromous fish. Although some mechanisms are becoming clear, many critical uncertainties remain.

Emigrating smolts are subject to intense predation as they enter estuaries (Järvi 1990, Dieperink et al. 2002) and predation rates may approach 25% (Hvidsten and Møkkelgjerd 1987, Hvidsten and Lund 1987). Thus many researchers consider smolt predation important in structuring salmon populations (Mather 1998). Smolts are especially vulnerable to predation upon entering saltwater because they encounter new habitats and predators while simultaneously being subject to osmotic stress while adapting to the saltwater environment (Järvi 1989). Stress levels alone may elevate predation risk to emigrating smolts by reducing the amount of energy available for normal activities (Barton and Schreck 1987) such as proper anti-predator behavior (Handeland et al. 1996). Thus the mechanism responsible for the observed high predation rates may be osmotic stress (Staurnes et al. 1996). This problem is likely exacerbated in many Maine rivers where the smoltification process appears to be especially difficult given the low pH experienced by emigrating smolts (Magee et al. 2001).

As more energy is required to simply maintain homeostatic balance, less energy is available for normal activities and predator avoidance behaviors appear to be compromised resulting in higher predation rates (Handeland et al. 1996). In addition, episodic acidification, such as those experienced during spring freshets (Whiting 2003), leads to disrupted osmoregulation as smolts transition to seawater (Magee et al. 2003) with mortality often occurring via predation in the wild (Kocik 2003 pers. comm.).

In addition to water quality perturbations, the decreased abundance of other anadromous stocks may also play a role in the perceived elevated predation risks as smolts and post-smolts transition to saltwater. Prior to European settlement, many anadromous species were more abundant than they are today. Many US and southern Canadian rivers historically supported large and diverse anadromous fish populations including alewife (*Alosa pseudoharengus*), blueback herring (*Alosa aestivalis*), American shad (*Alosa sapidissima*), and rainbow smelt (*Osmerus mordax*) in addition to Atlantic salmon. For some stocks, declines have been as dramatic as Atlantic salmon declines. For example, US landings of American shad exceeded 22,000 mt in 1896. Today, landings rarely exceed 2,000 mt and have averaged less than 1,350 mt since 1980 (Kocik 1998). These populations likely served as an important predation buffer allowing predators to focus on more abundant and energy-rich prey species (Schulze 1996, Brown et al. 2002). However, some of the same anthropogenic perturbations (e.g., dams, pollution, etc.) that lead to salmon declines have also diminished other anadromous populations (Collette and Klein-MacPhee 2002). The extent to which declines in other anadromous populations have led to higher smolt predation is yet to be assessed quantitatively.

### *Critical Uncertainties*

Sea surface temperature (SST) is another factor that may mediate Atlantic salmon predator-prey dynamics. Accumulating evidence suggests that changes in the SST regime in the northwest Atlantic can fundamentally change predator-prey dynamics. For example, Montevecchi et al. (2002) recorded a substantial increase in Atlantic salmon post-smolts in the diet of northern gannets beginning in 1990. This increase coincided with a shift from warm-water to cold-water pelagic prey (Montevecchi and Myers 1997). This shift was likely influenced by colder SSTs in the 1990s which likely inhibited highly migratory, warm-water mackerel (*Scomber scombrus*) and short-finned squid (*Illex illecebrosus*) from moving into the Newfoundland region (Templeman and Fleming 1953 cited in Montevecchi and Myers 1997). The extent to which changes in the SST regime are impacting other predator-prey relationships in the northwest Atlantic is largely unknown. The observations by Montevecchi and Meyers (1997) and Montevecchi et al. (2002) were made possible by long-term data sets that span the decadal changes in SST regimes. Thus similar patterns with other species may be occurring without our knowing because such comprehensive data sets are relatively rare.

Further uncertainty stems from the cumulative, perhaps synergistic (Järvi 1989), nature of the impacts of many anthropogenic perturbations to Atlantic salmon ecosystems. Järvi (1989) described the effects of osmotic stress and stress due to the presence of a predator as synergistic (i.e., greater than the sum of the two effects alone). Similarly, the impacts of dams, elevated predator abundance, and water quality perturbations may be leading to the very high mortality experienced by emigrating smolts in US rivers. For example, the presence of dams alone would likely have minimal impacts to emigrating smolts if the dams only delayed migration and killed a small portion of the total smolt biomass. However, when you consider the habitat preference of double-crested cormorants (Blackwell et al. 1997), the

migration delay due to dams (Blackwell et al. 1998), physical striking caused by turbines (Raymond 1988), decreased predator avoidance behaviors due to acid rain (Handeland et al. 1996), and the lack of emigrating Alosids as a predation buffer for emigrating smolts (Brown et al. 2002), a different picture emerges. Many anthropogenic impacts to Atlantic salmon ecosystems appear to cause system-wide destabilizations that are perceived as predator-prey dynamics issues.

Regardless of ultimate causes, many predation issues must be addressed as stocks of Atlantic salmon have been listed as endangered in the US and Canada. There are several approaches to consider as we move forward with Atlantic salmon research and management. Further research is clearly warranted as long as the problems with quantifying marine predation, clearly outlined by Cairns (2001a), are considered. The results of further studies may be problematic for US managers as critically low abundance further exacerbates the problem of even documenting predation in the open ocean. However, Cairns' (2001a) proposed assessments of life history, ocean temperatures, and schooling behavior would clearly provide important insights into the tradeoffs between predation risk and high growth potential. Further examination of novel approaches (e.g., fatty acid signature analysis, genetic analysis of seal scat) to detection and measurement of predation may also produce useful information for identifying localized predation issues.

### **Management Measures and On-going Research**

Several management measures aimed at reducing early marine predation have recently been initiated in the US. First, an experimental liming project has been initiated. The objective of the project is to ease the transition from freshwater to saltwater for emigrating smolts in the Dennys River in eastern Maine. This approach might ameliorate the impacts of episodic low pH events (including low calcium and high aluminum concentrations) on the smoltification process, hopefully leading to higher early marine survival. If successful, this technique may be expanded to other salmon rivers that experience episodic low pH events. Second, an experimental double-crested cormorant control project has been initiated. This project is using non-lethal methods displace foraging double-crested cormorants from the lower Narraguagus River and estuary during the smolt migration. The goal is to see if displacement alone can have a measurable effect upon smolt survival as they enter the ocean. If successful, this programme may be expanded in the near future. If unsuccessful, the project may also use lethal removals in subsequent years. These are both adaptive management experiments that allow managers to evaluate the effects of their actions and adjust strategies accordingly. No results are yet available from either study.

In addition, on-going research at the University of Maine is attempting to document the frequency, pattern, and extent of seal depredations at salmon aquaculture farms. This research should assist the aquaculture industry by identifying the most successful mitigation measures available in US waters. This research could also assist wild salmon recovery efforts by minimizing domesticated salmon escapement.

Several other management options that are yet to be tested in the US are also available. Non-lethal seal exclusion from a particular estuary could be examined. This approach has recently been shown to deter harbor seal predation on Pacific salmon in British Columbia, Canada (Yurk and Trites 2000) and these techniques may be transferable. Second, open sea releases of hatchery smolts could be used to minimize predation as smolts transition to seawater. This approach increased return rates by as much as 111% in Norway (Gunnerød et al. 1988).

These additional management options may seem somewhat drastic but may be worth pursuing given the endangered status of some US stocks. Each technique would also require an adaptive management approach.

## References

- Amiro, P. G. 1998. The abundance of Harp seals in the North Atlantic and recruitment of the North American stock of Atlantic salmon (*Salmo salar*). Canadian Stock Assessment Secretariat Research Document 98/84.
- Anthony, V. C. 1994. The Significance of Predation on Atlantic Salmon. *In*. A Hard Look at Some Tough Issues: Proceedings of the New England Atlantic Salmon Management conference. Edited by S. Calabi and A. Stout. New England Salmon Association. Newburyport, Massachusetts. pp 240-284.
- Anthony, V.C. 1996. The predators of Atlantic salmon and their impact on salmon stocks. Pages 5-65 in "The Atlantic salmon as predator and prey". North Atlantic Salmon Conservation Organization. Edinburgh, Scotland.
- Barton, B. A. and C.B. Schreck. 1987. Influence of acclimation temperature on interrenal and carbohydrate stress responses in juvenile chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 62: 299-310.
- Baum, E. 1997. Maine Atlantic Salmon: a national treasure. Atlantic Salmon Unlimited. Hermon, Maine.
- Beland, K.F., J.F. Kocik, J. vandeSande, and T.F. Sheehan. 2001. Striped bass predation upon Atlantic salmon smolts in Maine. *Northeastern Naturalist* 8(3): 267-274.
- Blackwell, B.F. and F. Juanes. 1998. Predation on Atlantic salmon smolts by striped bass after dam passage. *North American Journal of Fisheries Management* 18: 936-939.
- Blackwell, B.F., W.B. Krohn, N.R. Dube, and A.J. Godin. 1997. Spring prey use by double-crested cormorants on the Penobscot River, Maine, USA. *Colonial Waterbirds* 20(1): 77-86.
- Blackwell, B.F., and W.B. Krohn. 1997. Spring foraging distribution and habitat selection by Double-crested Cormorants on the Penobscot River, Maine. *Colonial Waterbirds* 20:66-76
- Blackwell, B.F., G. Gries, F. Jauanes, K.D. Friedland, L.W. Stolte and J.F. McKeon. 1998. Simulating migration mortality of Atlantic salmon smolts in the Merrimack River. *North American Journal of Fisheries Management* 18:31-45.
- Boyle, P.R., G.J. Pierce, and J.S.W. Diack. 1990. Sources of evidence for salmon in the diet of seals. *Fisheries Research* 10: 137-150.
- Brown, R.W., F. Juanes, and G. Shepard. 2002. Approaches to evaluating the role of marine fish predation on juvenile Atlantic salmon. Sixth International Atlantic Salmon Symposium. Edinburgh, Scotland.



- Cairns, D.K. 1998. Diet of cormorants, mergansers, and kingfishers in northeastern North America. Canadian Technical Report of Fisheries and Aquatic Sciences No. 2225.
- Cairns, D.K. 2001a. Approaches and methods for the scientific evaluation of bird and mammal predation on salmon in the Northwest Atlantic. Canadian Stock Assessment Secretariat Research Document 2001/011.
- Cairns, D. K. 2001b. An evaluation of possible causes of the decline in pre-fishery abundance of North American Atlantic salmon. Canadian Technical Report of Fisheries and Aquatic Sciences No. 2358.
- Cairns, D. K. and D.G. Reddin. 2000. The potential impact of seal and seabird predation on North American Atlantic salmon. Canadian Stock Assessment Secretariat Research Document 2000/12.
- Clifford, S. L., P. McGinnity, and A. Ferguson. 1998. Genetic changes in Atlantic salmon (*Salmo salar*) populations of northwest Irish rivers resulting from escapes of adult farm salmon. Canadian Journal of Fisheries and Aquatic Sciences 55: 358-363.
- Collette, B. B. and G. Klein-MacPhee. 2002. Fishes of the Gulf of Maine, Third Edition. Smithsonian Institution Press. Washington. D.C.
- Diepernick, C., B.D. Bak, L.F. Pedersen, M.I. Pedersen, and S. Pedersen. 2002. Predation on Atlantic salmon and sea trout during their first days as postsmolts. Journal of Fish Biology 61: 848-852.
- Field, J.D. 1997. Atlantic striped bass management: Where did we go right? Fisheries 22(7):6-8.
- Fiscus, C.H. 1980. Marine mammal-salmonid interactions: a review. In Salmonid Ecosystems of the North Pacific. Edited by W.J. McNeil and D.C. Himsworth Oregon State University, Corvallis. pp. 121-132.
- Gilbert, J.R. 2003. University of Maine, Professor of Wildlife Ecology. Personal Communication.
- Gulland, J.A. 1987. Seals and fisheries: A case for predator control? Trends in Ecology and Evolution 2(4): 102-104.
- Gunnerød, T. B., N.A. Hvidsten, and T.G. Heggberget. 1988. Open sea releases of Atlantic salmon smolts, *Salmo salar*, in central Norway, 1973-83. Canadian Journal of Fisheries and Aquatic Sciences 45: 1340-1345.
- Hammill, M.O., and G.B. Stenson. 2000. Estimated prey consumption by harp seal (*Phoca groenlandica*), hooded seals (*Cystophora cristata*), grey seals (*Halichoerus grypus*) and harbour seals (*Phoca vitulina*) in Atlantic Canada. Journal of Northwest Atlantic Fishery Science 26: 1-23.

- Handelhand, S.O., T. Järvi, A. Fernö, and S.O. Stefansson. 1996. Osmotic stress, antipredator behavior, and mortality of Atlantic salmon (*Salmo salar*) smolts. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 2673-2680.
- Harwood, J. 1984. Seals and Fisheries. *Marine Pollution Bulletin* 15(12): 426-429.
- Hvidsten, N. A. and R.A. Lund. 1988. Predation on hatchery-reared and wild smolts of Atlantic salmon, *Salmo salar* L., in the estuary of River Orkla, Norway. *Journal of Fish Biology* 33: 121-126.
- Hvidsten, N.A. and P.I. Møkkelgjerd. 1987. Predation on salmon smolts, *Salmon salar* L., in the estuary of the River Surna, Norway. *Journal of Fish Biology* 30: 273- 280.
- Järvi, T. 1989. Synergistic effect on mortality in Atlantic salmon, *Salmo salar*, smolt caused by osmotic stress and presence of predators. *Environmental Biology of Fishes* 26: 149-152.
- Järvi, T. 1990. Cumulative acute physiological stress in Atlantic salmon smolts: the effect of osmotic imbalance and the presence of predators. *Aquaculture* 89:337-350.
- Kocik J.F. 1998. American shad. *In* Status of fishery resources off the Northeastern United States for 1998. *Edited by* S.H. Clark. NOAA Technical Memo NMFS-NE-115. pp. 136-137.
- Kocik, J.F. 2003. Research Fishery Biologist, National Marine Fisheries Service. Personal Communication.
- Larsson, P.O. 1985. Predation on migrating smolt as a regulating factor in Baltic salmon, *Salmo salar* L., populations. *Journal of Fish Biology* 26: 391-397.
- Lawson, J.W., J.T. Anderson, E.L. Dalley, and G.B. Stenson. 1998. Selective foraging by harp seals *Phoca groenlandica* in nearshore and offshore waters of Newfoundland, 1993 and 1994. *Marine Ecology Progress Series* 163: 1-10.
- Magee, J.A., T.A. Haines, J.F. Kocik, K.F. Beland, and S.D. McCormick. 2001. Effects of acidity and aluminum on the physiology and migratory behavior of Atlantic salmon smolts in Maine, USA. *Water, Air, and Soil Pollution* 130: 881-886.
- Magee, J.A., M. Obedzinski, S.D. McCormick, and J.F. Kocik. 2003. Effects of episodic acidification on Atlantic salmon (*Salmo salar*) smolts. *Canadian Journal of Fisheries and Aquatic Sciences* 60: 214-221.
- Mather, M.E. 1998. The role of context-specific predation in understanding patterns exhibited by anadromous salmon. *Canadian Journal of Fisheries and Aquatic Sciences* 55(Suppl. 1): 232-246.
- Meister, A. L. 1984. The marine migrations of tagged Atlantic salmon (*Salmo salar* L.) of USA origin. Anadromous and Catadromous Fish Committee, International Council for the Exploration of the Sea. Copenhagen, Denmark.

- Meister, A. L. and F.J. Gramlich. 1967. Cormorant predation on tagged Atlantic salmon smolts. Final report of the 1966-67 Cormorant - Salmon Smolt Study. Atlantic Sea Run Salmon Commission. Orono, Maine.
- Montevecchi, W.A., D.K. Cairns, and R.A. Myers. 2002. Predation on marine-phase Atlantic salmon (*Salmo salar*) by gannets (*Morus bassanus*) in the Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 602-612.
- Montevecchi, W.A. and R. A. Myers. 1997. Centurial and decadal oceanographic influences on changes in northern gannet populations and diets in the north-west Atlantic: implications for climate change. *ICES Journal of Marine Science* 54: 608-614.
- Moring, J.R., O. van den Ende, and K.S. Hockett. 1999. Predation on Atlantic salmon smolts in New England rivers. *In Smolt, physiology, ecology and behaviour. Edited by S. McCormick and D. MacKinlay. Department of Fisheries and Oceans, Vancouver, British Columbia, Canada. pp. 127-139.*
- Pierce, G. J., P.M. Thompson, A. Miller, J.S.W. Diack, D. Miller, and P.R. Boyle. 1991. Seasonal variation in the diet of common seals (*Phoca vitulina*) in the Moray Firth area of Scotland. *Journal of Zoology* 223: 641-652.
- Raymond, H.L. 1988. Effects of Hydroelectric Development and Fisheries Enhancement on Spring and Summer Chinook Salmon and Steelhead in the Columbia River Basin. *North American Journal of Fisheries and Aquatic Sciences* 8: 1-24.
- Saunders, R. L. 1991. Potential interactions between cultured and wild Atlantic salmon. *Aquaculture* 98: 51-60.
- Schulze, M.B. 1996. Using a field survey to assess potential temporal and spatial overlap between piscivores and their prey, and a bioenergetics model to examine potential consumption of prey, especially juvenile anadromous fish in the Connecticut River estuary. M.Sc. Thesis. University of Massachusetts, Amherst. 133 pp.
- Staurnes, M., L.P. Hansen, K. Fugelli, and O. Haraldstad. 1996. Short-term exposure to acid water impairs osmoregulation, seawater tolerance, and subsequent marine survival of smolts of Atlantic salmon (*Salmo salar* L.). *Canadian Journal of Fisheries and Aquatic Sciences* 53: 1695-1704.
- Thompson, P. M. and F. Mackay. 1999. Pattern and prevalence of predator damage on adult Atlantic salmon, *Salmo salar* L., returning to a river system in north-east Scotland. *Fisheries Management and Ecology* 6: 335-343.
- Templeman, W. and A.M. Fleming. 1953. Long-term changes in hydrographic conditions and corresponding changes in the abundance of marine animals. *Canadian Journal of Fisheries and Aquatic Sciences* 42: 976-981.
- Van den Ende, O. 1993. Predation on Atlantic salmon smolts (*Salmo salar*) by smallmouth bass (*Micropterus dolomeiu*) and chain pickerel (*Esox niger*) in the Penobscot River, Maine. M.Sc. Thesis. University of Maine, Orono. 95 pp.

- Webb, J. H., A.F. Youngson, C.E. Thompson, D.W. Hay, M.J. Donagy, and I.S. McLaren. 1983. Spawning of escaped farmed Atlantic salmon, *Salmo salar* L., in western and northern Scottish rivers: egg deposition by females. *Aquaculture and Fisheries Management* 24: 663-670.
- Whiting, M. 2003. Maine salmon rivers water quality monitoring progress report for the 2002 field season. Maine Department of Environmental Protection, Bangor, Maine, 04401.
- Windsor, M. L. and P. Hutchinson. 1990. The potential interactions between salmon aquaculture and the wild stocks - a review. *Fisheries Research* 10: 163-176.
- Youngson, A. F., J.H. Webb, C.E. Thompson, and D. Knox. 1993. Spawning of escaped farmed Atlantic salmon (*Salmo salar*): hybridization of females with brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* 50: 1986-1990.
- Yurk H., and A.W. Trites. 2000. Experimental Attempts to Reduce Predation by Harbor Seals on Out-Migrating Juvenile Salmonids. *Transactions of the American Fisheries Society* 129: 1360-1366.